

3-dimensional low-energy topological invariants

Małgorzata Bakalarska and Bogusław Broda

Department of Theoretical Physics, University of Łódź
 Pomorska 149/153, PL-90-236 Łódź, Poland
 gosiabak@kryisia.uni.lodz.pl
 bobroda@kryisia.uni.lodz.pl

Abstract: A description of the one-loop approximation formula for the partition function of a three-dimensional abelian version of the Donaldson-Witten theory is proposed. The one-loop expression is shown to contain such topological invariants of a three-dimensional manifold $\mathcal{M}$ like the Reidemeister-Ray-Singer torsion τ_R and Betti numbers.

1. In three dimensions (3d), traditionally, we have the two important well-known topological quantum field theories (of “cohomological” type): non-abelian $SU(2)$ topological gauge theory of flat connection and 3d version of the topological Seiberg-Witten theory. The former is a 3d twisted $\mathcal{N} = 4$ SUSY $SU(2)$ pure gauge theory or a 3d version of the Donaldson-Witten theory, and “by definition” it describes the Casson invariant, which appropriately (i.e. with sign) counts the number of flat $SU(2)$ connections [1]. The latter is a dimensional reduction of the topological Seiberg-Witten theory or a twisted version of 3d $\mathcal{N} = 4$ SUSY $U(1)$ gauge theory with a matter hypermultiplet [2], [3]. Also this theory describes an interesting non-trivial topological invariant of 3d manifolds pertaining to 3d Donaldson-Witten invariant which has been conjectured to be equivalent to already known topological invariants and in particular to topological (Reidemeister) torsion [4]. The conjectures are physically strongly motivated by the fact that the both theories can be derived, performing dimensional reduction and/or taking low-energy limit, from 4d $\mathcal{N} = 2$ SUSY $SU(2)$ pure gauge theory [5] corresponding via twist to Donaldson-Witten theory. Interestingly, it follows from [6] that low-energy effective theory for both 3d $\mathcal{N} = 4$ SUSY $SU(2)$ pure gauge case as well as for the $\mathcal{N} = 4$ SUSY abelian one with matter hypermultiplet is an abelian pure gauge theory. Therefore, we should essentially deal with 3d twisted version of the abelian theory—topological gauge theory of flat $U(1)$ -connection [7]. In dual variables, we have a 3d $\mathcal{N} = 4$ SUSY σ -model with a hyper-kähler manifold as a target space. Twisted version of this model has been analyzed in detail by Rozansky and Witten in [8]. Here, we will consider a corresponding topological model in original (non-dual) variables. In fact, we will confine ourselves to the lowest-order one-loop calculation which nevertheless gives a (exact) topological invariant—Reidemeister-Ray-Singer torsion—as it should. In spite of a direct interpretation of our model as a low-energy version of the above-mentioned gauge models, we can treat it as a stand-alone one as well. Furthermore, we can even treat its (one-loop) gaussian approximation as an independent model, as we actually do in this note.

2. Topological action of our theory assumes the following form

$$S_{\text{eucl}} = \int_{\mathcal{M}} d^3x \sqrt{g} \left\{ \frac{1}{4} F^{mn} F_{mn} - \frac{1}{2} \varphi \nabla^m \nabla_m \varphi - \bar{B} \nabla^m \nabla_m B - \frac{1}{\sqrt{g}} \varepsilon^{kmn} \chi_k \partial_m \psi_n + \right. \\ \left. - \omega \nabla^m \chi_m + \rho \nabla^m \psi_m \right\}, \quad (1)$$

where F_{mn} is the usual $U(1)$ -gauge field strength, B is a complex scalar, φ is a real scalar, and the fields ω , ρ , χ_k , ψ_k are fermionic fields. The covariant derivative of fermions and bosons, here denoted as ∇_m , is defined using the Levi-Civita connection on $\mathcal{M}$ with a metric g_{mn} , where $m, n, k = 1, 2, 3$.

Gauge invariance of the action (1) requires gauge fixing and introduction of the Faddeev-Popov ghost fields c , $\bar{c}$

$$\mathcal{L}_{\text{gauge}} = \frac{1}{2} (\nabla^m A_m)^2 - \bar{c} \nabla^m \nabla_m c. \quad (2)$$

Supplementing the action (1) with (2), we obtain

$$S_{\text{eucl}} = \int_{\mathcal{M}} d^3x \sqrt{g} \left\{ \frac{1}{4} F^{mn} F_{mn} - \frac{1}{2} \varphi \nabla^m \nabla_m \varphi - \bar{B} \nabla^m \nabla_m B + \right. \\ \left. - \frac{1}{\sqrt{g}} \varepsilon^{kmn} \chi_k \partial_m \psi_n - \omega \nabla^m \chi_m + \rho \nabla^m \psi_m + \right. \\ \left. + \frac{1}{2} (\nabla^m A_m)^2 - \bar{c} \nabla^m \nabla_m c \right\}. \quad (3)$$

It will appear that the partition function Z corresponding to (3) essentially consists of the Reidemeister-Ray-Singer torsion of 3d manifold $\mathcal{M}$. Namely,

$$Z(\mathcal{M}) = \int [\mathcal{D}X] \exp \{ -S_{\text{eucl}}[X] \}, \quad (4)$$

where the integration measure $[\mathcal{D}X]$ is taken over all the fields

$$(A_m, \varphi, B, \bar{B}, \omega, \rho, \chi_k, \psi_k, c, \bar{c}).$$

The path integral over the bosonic field φ gives the factor

$$(\det'(-\Delta_0))^{-\frac{1}{2}}, \quad (5)$$

where $\Delta_0 = \nabla^m \nabla_m$ is a Laplacian acting on zero-forms on $\mathcal{M}$ and $\det'$ means that we exclude zero modes. The integration over the usual ghosts c and $\bar{c}$ gives $\det'(-\Delta_0)$, while the B - and $\bar{B}$ -integrations give its inverse $(\det'(-\Delta_0))^{-1}$. So, these modes are canceled. Now let us introduce an operator L_- which acts on the direct sum of zero- and one-forms on $\mathcal{M}$ [8]

$$L_-(\rho, \chi_k) = (-\nabla^m \chi_m, \nabla^k \rho + \frac{1}{\sqrt{g}} \varepsilon^{kmn} \partial_m \chi_n), \quad (6)$$

and similarly for (ω, ψ_k) .

Following [8] we define a scalar product

$$\langle \rho, \chi_k | \rho', \chi'_k \rangle = \int_{\mathcal{M}} d^3x \sqrt{g} (\rho \rho' + \chi^k \chi'_k). \quad (7)$$

Then the part of the action (3) containing ω , ρ , χ_k and ψ_k becomes a quadratic form

$$\frac{1}{2} \langle \omega, \psi_k | L_- | \rho, \chi_k \rangle - \frac{1}{2} \langle \rho, \chi_k | L_- | \omega, \psi_k \rangle. \quad (8)$$

The fermionic one-loop contribution [9], with zero modes removed, is

$$\det' L_- . \quad (9)$$

Finally, the integration over the gauge field A_m gives

$$(\det'(-\Delta_1))^{-\frac{1}{2}}, \quad (10)$$

so that the total one-loop contribution of non-zero modes is

$$\frac{\det' L_-}{(\det'(-\Delta_0))^{\frac{1}{2}} (\det'(-\Delta_1))^{\frac{1}{2}}} . \quad (11)$$

The absolute value of the ratio of the determinants in eq.(11) is related to the Reidemeister-Ray-Singer analytic torsion $\tau_R(\mathcal{M})$ [8], [10], [11]. The formula for the absolute value of the ratio of the determinants is

$$\left| \frac{\det' L_-}{(\det'(-\Delta_0))^{\frac{1}{2}} (\det'(-\Delta_1))^{\frac{1}{2}}} \right| = \tau_R^{-2}(\mathcal{M}). \quad (12)$$

3. It is not the whole story, because the partition function is, in principle, plagued by zero modes, which we have removed by hand. There are the following four sorts of zero modes, which should be treated differently (the number of some of them may depend on topology of $\mathcal{M}$): (1) 3 scalar boson zero modes, corresponding to φ , B and $\bar{B}$; (2) b_1 ($= 1^{\text{st}}$ Betti number of $\mathcal{M}$) vector boson zero modes of A_m ; (3) 1 ghost zero mode for c , $\bar{c}$; (4) 2 scalar fermion zero modes of ω , ρ , and $2b_1$ one-form zero modes for χ_k , ψ_k . The ghost zero mode can be removed instantaneously because it should not be present in the partition function from the very beginning at all, as the gauge transformation corresponding to the constant (zero) mode acts trivially on A_m . Possibly, we could integrate out eq.(4) with respect to vector zero modes because it corresponds to integrating over b_1 -torus. The rest of boson and fermion zero modes produces infinities and zeros in the partition function respectively. In this gaussian approximation/version there is no other possibility than to remove them by hand. They can be properly handled only in higher orders. We need a compact-space valued scalar field (σ -model) to integrate out scalars, and higher-order fermion terms to saturate fermion zero modes. As usual, boson and ghost zero modes yield appropriate powers in the Planck constant in front of eq.(4).

The second author (B. B.) would like to thank the organizers for their kind invitation and assistance during the symposium. The work has been supported by the KBN grant 2 P03B 084 15.

References

- [1] M. Blau, G. Thompson, N=2 topological gauge theory, the Euler characteristic of moduli spaces, and the Casson invariant, *Commun. Math. Phys.* **152**, 41(1993).
- [2] N. Seiberg, E. Witten, Electric-magnetic duality, monopole condensation, and confinement in N=2 supersymmetric Yang-Mills theory, *Nucl. Phys.* **B426**, 19(1994).
- [3] E. Witten, Monopoles and four-manifolds, *Math. Res. Lett.* **1**, 769(1994).
- [4] B. Broda, Topological contents of 3D Seiberg-Witten theory, *New Developments in Quantum Field Theory* (NATO ASI Series, Zakopane, June 1997, ed. P.H. Damgaard), Plenum Press, pp. 261–268.
- [5] E. Witten, Topological quantum field theory, *Commun. Math. Phys.* **117**, 353(1988).
- [6] N. Seiberg, E. Witten, Gauge Dynamics And Compactification To Three Dimensions, *E-print* hep-th/9607163.
- [7] M. Bakalarska, B. Broda, 3-dimensional topological quantum field theory of Witten type, Proceedings of the International Conference “*Particles, Fields and Gravitation*”, Lodz, April 1998, ed. J.Rembieliński, American Institute of Physics, Woodbury, New York, 1998, pp. 303–311.
- [8] L. Rozansky, E. Witten, Hyper-Kähler geometry and invariants of three-manifolds, *E-print* hep-th/9612216.
- [9] E. Witten, Quantum field theory and the Jones polynomial, *Commun. Math. Phys.* **121**, 351(1989).
- [10] L. Rozansky, A large k asymptotics of Witten’s invariant of Seifert manifolds, *Commun. Math. Phys.* **171**, 279(1995).
- [11] L. Rozansky, Witten’s invariant of 3-dimensional manifolds: loop expansion and surgery calculus, *Knots and Applications*, ed. L. H. Kauffman, 271–299.